

This Information Sheet describes the *typical average properties* of the specified soil. It is essentially a summary of information obtained from one or more profiles of this soil that were examined and described during the Topoclimate survey or previous surveys. It has been prepared in good faith by trained staff within time and budgetary limits. However, no responsibility or liability can be taken for the accuracy of the information and interpretations. Advice should be sought from soil and landuse experts before making landuse decisions on individual farms and paddocks. The characteristics of the soil at a specific location may differ in some details from those described here.
No warranties are expressed or implied unless stated.

Soil name: **Crookston**

Overview

Crookston soils occupy about 6,100 ha on terraces in the Waimea plains, and in fans flanking the Blue Mountains between Heriot and Tapanui. They are formed in wind deposited loess derived from greywacke and schist rocks. Crookston soils are well drained and have a deep rooting depth, high water holding capacity, and have light silt loam textures with P-retention between 20 and 40%. They are high producing soils currently used for intensive sheep and dairy production and some cropping. Rainfall is evenly spread, although these soils can be seasonally dry over the summer.

Physical properties

Crookston soils have a deep rooting depth and high plant available water, meaning there is no significant physical barrier to root growth. The soils are well drained but the compact subsoil is slowly permeable, and may cause short-term waterlogging after heavy rainfall. Texture is light silt loam in all horizons, with topsoil clay content of 20–25%. Crookston soils are typically stone free, although the moderately deep phases have gravels between 45 and 90cm depth that may restrict rooting depth and reduce available water status to moderately high.



Crookston profile

Fertility properties

Topsoil organic matter values range from 5 to 7%, P-retention values 20–40% and pH values moderate. Cation exchange and base saturation values are high in the topsoil and decrease down the profile. Available calcium, magnesium and potassium values are moderate to high in the upper soil horizons. Phosphorus reserves are low and sulphur levels low in the topsoil but high in the subsoil. Micro-nutrient levels are generally adequate although boron responses in brassicas and molybdenum responses in legumes can occur.

Associated and similar soils

Some soils that commonly occur in association with Crookston soils are:

- Kaweku: well drained, shallow soils on high terraces
- Arthurton: imperfectly drained equivalent of the Crookston soil
- Waikoikoi: moderately deep to deep; poorly drained due to fragipan
- Jacobstown: poorly drained floodplain soil due to a high groundwater table

Some soils that have similar properties to Crookston soils are:

- Clinton: similar soil showing a greater degree of Brown soil properties with browner colours and P-retention of 30–40% throughout the profile; occurs on fans and terraces near Clinton
- Tuturau: similar soil but has loamy silt subsoil textures; formed in near-source loess adjacent to the Mataura River, between Mataura and Waimahaka
- Waikaka: also has Brown-Pallic intergrade properties, but has an upper subsoil with browner colours and P-retention of 40–60%; occurs in higher rainfall rolling and hilly land grading between the downlands and the hill country
- Nokomai: dominated by Pallic soil properties; occurs in the Garston and Waikaia plains areas
- Edendale: Brown soil on intermediate terraces of the Southland plain; has P-retention of 60–80%

Sustainable management indicators

Note: the vulnerability ratings given in the table below are generalised and should not be taken as absolutes for this soil type in all situations. The actual risk depends on the environmental and management conditions prevailing at a particular place and time. Specialist advice should be sought before making management decisions that may have environmental impacts. Where vulnerability ratings of Moderate to Very severe are indicated, advice may be sought from Environment Southland or a farm management consultant.

Vulnerability factor	Rating	Vulnerability compared to other Southland soils
Structural compaction	moderate	These soils have a moderate vulnerability to structural degradation by long-term cultivation, or compaction by heavy stocking and vehicles. This rating reflects the light silt loam texture and low P-retention.
Nutrient leaching	moderate	These soils have a moderate vulnerability to leaching to groundwater. This rating reflects the high water-holding capacity and slow subsoil permeability, but is offset by the good profile drainage.
Topsoil erodibility by water	moderate	Due to the light silt loam texture, the topsoil erodibility of these soils is moderate. Erodibility is highly dependent on management, particularly when there is no vegetation cover.
Organic matter loss	slight	Vulnerability to long-term decline in soil organic matter levels is partly dependent on soil properties, and highly dependent on management practices (e.g., crop residue management and cultivation practices).
Waterlogging	slight	These soils have a slight vulnerability to waterlogging during wet periods. This rating reflects the good drainage, but slowly permeable subsoil.

General landuse versatility ratings

Note: The versatility ratings in the table below are indicative of the major limitations for semi-intensive to intensive land use. These ratings differ from those used in the past in that sustainability factors are incorporated in the classification. Refer to the Topoclimate district soil map or property soil map to determine which of the soil symbols listed below are applicable, then check the versatility ratings for that symbol in the appropriate table.

CkU1 (Crookston undulating deep)

Versatility evaluation for soil CkU1

Landuse	Versatility rating	Main limitation
Non-arable horticulture	Moderate	Risk of short-term waterlogging after heavy rain
Arable	Moderate	Risk of short term waterlogging after heavy rain
Intensive pasture	Moderate	Vulnerability to nutrient leaching to ground water; vulnerability to structural compaction.
Forestry	High	Few limitations

CkU2 (Crookston undulating moderately deep)

Versatility evaluation for soil CkU2

Landuse	Versatility rating	Main limitation
Non-arable horticulture	Moderate	Risk of short term waterlogging after heavy rain; restricted rooting depth.
Arable	Moderate	Risk of short- term waterlogging after heavy rain
Intensive pasture	Moderate	Vulnerability to nutrient leaching to ground water; vulnerability to structural compaction.
Forestry	Moderate	Restricted rooting depth.

CkR1 (Crookston rolling deep)

Versatility evaluation for soil CkR1

Landuse	Versatility rating	Main limitation
Non-arable horticulture	Moderate	Risk of short-term waterlogging after heavy rain; rolling slopes
Arable	Limited	Rolling slopes
Intensive pasture	Moderate	Rolling slopes; vulnerability to structural compaction
Forestry	High	Few limitations

Management practices that may improve soil versatility

- Careful management after heavy rainfall and wet periods will reduce the impact of short-term waterlogging. Intensive stocking, cultivation and vehicular traffic should be minimised during these periods.
- Installation and maintenance of subsurface mole and tile drainage will reduce the risk of short-term waterlogging
- Management of nutrient applications to minimise leaching losses.

Copyright © 2002, Crops for Southland

www.cropssouthland.co.nz

This Information Sheet may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgement of the source is made. Crops for Southland and Environment Southland would appreciate receiving a copy of any publication that uses this Information Sheet as a source. No use of this Information Sheet may be made for resale or for any other commercial purpose whatsoever without prior permission in writing from Crops for Southland.