

This Technical Data Sheet describes the *typical average properties* of the specified soil.

It is essentially a summary of information obtained from one or more profiles of this soil that were examined and described during the Topoclimate survey or previous surveys. It has been prepared in good faith by trained staff within time and budgetary limits. However, no responsibility or liability can be taken for the accuracy of the information and interpretations. Advise should be sought from soil and landuse experts before making landuse decisions on individual farms and paddocks. The characteristics of the soil at a specific location may differ in some details from those described here.

No warranties are expressed or implied unless stated.

Soil name: **McKerchar**

Overview

McKerchar soils occupy about 100 ha on floodplains of streams from the Takitimu Mountains, mainly between the Blackmount and The Key areas. These soils also occur on areas outside the Topoclimate survey. They are formed into shallow fine alluvium over gravels that are derived from tuffaceous greywacke and basic volcanic rocks. Soils are shallow, with clayey textures, and poorly drained due to a high water-table. Present use is pastoral grazing with sheep and beef cattle. Climate is temperate with regular rainfall throughout the year. Soils are rarely dry.

Soil classification

NZ Soil Classification (NZSC):

Typic Recent Gley; rounded-stony, tuffaceous sandstone; clayey.

Previous NZ Genetic Classification:

Recent soils

Classification explanation

McKerchar soils have been reclassified in this survey because the poor drainage correlates to gley soils rather than the previous classification. The soils are poorly drained due to a high groundwater table, and accumulation of sediment is sufficient that subsoils show little structural development. They have silty clay textures and are shallow, with less than 45cm to the gravels.

Soil phases and variants

Identified units in the McKerchar soils are:

- McKerchar undulating shallow (McU3): has gravel within 45cm depth; occurs on slopes of 0–7°

The soil properties described in this Technical Data Sheet are based on the most common phase, McKerchar undulating shallow (McU3). Values for other phases and variants can be taken as being similar. Where they differ significantly they are recorded with a separate versatility rating.

Associated soils

Some soils that commonly occur in association with McKerchar soils are:

- Waiau: well drained shallow soil on the floodplain and low terraces, with gravels within 45cm depth
- Tuatapere: well drained moderately deep to deep soil on the floodplain with gravels below 45cm
- Upukerora: soils of the active floodplain and riverbed; dominantly well drained and stony.


Similar soils

Some soils that have similar properties to McKerchar soils are:

- McLeish: occurs on low terraces of the Aparima River in the Drummond district; is slowly to non-accumulating, and subsoils show structural development
- Lumsden: occurs on floodplains of major streams and rivers in northern and central Southland and in west Otago; has silty textures
- Otepunui: forming predominantly into quartz gravels on stream floodplains of the Southland Plain

Typical profile features

The following is a 'generic' or composite profile description representing the most common combination of characteristics for this soil type. The actual profiles for which descriptions and data are available are listed at the end of this Technical Data Sheet.

McKerchar profile	Horizon	Depth (cm)	Description
	Apg	0–24	Brownish black slightly gravelly silty clay; few dull brown and very dark reddish brown mottles; weak soil strength; weakly developed fine polyhedral structure; abundant roots.
	BCg	24–44	Greyish yellow-brown slightly gravelly silty clay; many dull yellowish brown mottles; weak soil strength; massive structure; many roots.
	2Cg	44–65+	Extremely gravelly light grey sand; dense particle packing; single grain structure; gravels rounded and slightly weathered; no roots.

Key profile features

McKerchar topsoils are about 15–25cm deep with a weak structure. Subsoils are generally structureless to the underlying gravels. The dominance of grey colours throughout the subsoil reflects the poor drainage of the soils.

Typical physical properties

Note: values in *Italics* are estimates

Horizon	Depth (cm)	Bulk density	Permeability	Texture	Gravel content
Apg	0–24	—	<i>Moderate</i>	Silty clay	Slightly gravelly
BCg	24–44	—	<i>Moderate</i>	Silty clay	Slightly gravelly
2Cg	44–65+	—	<i>Slow</i>	Sand	Extremely gravelly

Profile drainage: Poor
Plant readily available water: *Moderate*
Potential rooting depth: Slightly deep
Rooting restriction: Gravelly subsoils

Key physical properties

McKerchar soils have a slightly deep rooting depth (45–60cm) with moderate plant available water. The soils are poorly drained, with moderate to slow permeability, and a high water table that is likely to cause significant aeration limitations during wet periods. Textures are silty clay with topsoil clay content about 35–45%. Soils have gravel in all horizons, and are very gravelly within 45cm depth.

Typical chemical properties

Horizon	Depth (cm)	pH	P retention	CEC	BS	Ca	Mg	K	Na
Apg	0–24	Moderate	High	Very high	Very low	Low	Very low	Low	Very low
BCg	24–44	Moderate	Very high	High	Very low	Low	Very low	Very low	Very low
2Cg	44–65+	—	—	—	—	—	—	—	—

Key chemical properties

Topsoil organic matter content is about 15–20%; P-retention >80% and pH moderate (mid 5s). Cation exchange is very high and base saturation very low. Available calcium, magnesium and potassium values are all low. Soil reserve phosphorus levels are low. Micronutrient levels are generally adequate.

Vulnerability to environmental degradation

Note: the vulnerability ratings given in the table below are generalised and should not be taken as absolutes for this soil type in all situations. The actual risk depends on the environmental and management conditions prevailing at a particular place and time. Specialist advice should be sought before making management decisions that may have environmental impacts. Where vulnerability ratings of Moderate to Very severe are indicated, advice may be sought from Environment Southland or a farm management consultant.

Vulnerability factor	Rating	Vulnerability compared to other Southland soils
Structural compaction	slight	These soils have a slight vulnerability to structural degradation by long-term cultivation, or compaction by heavy stocking and vehicles. This rating reflects the poor drainage, but strongly offset by the high clay, organic matter, and P-retention values.
Nutrient leaching	moderate	These soils have a moderate vulnerability to leaching to groundwater. This rating reflects the poor drainage and slow permeability, but moderate water-holding capacity.
Topsoil erodibility by water	minimal	Due to the high clay and organic matter content, topsoil erodibility in these soils is minimal. Erodibility is highly dependent on management, particularly when there is no vegetation cover.
Organic matter loss	moderate	Vulnerability to long-term decline in soil organic matter levels is partly dependent on soil properties and highly dependent on management practices (e.g., crop residue management and cultivation practices).
Waterlogging	severe	These soils have a severe vulnerability to waterlogging during wet periods. This rating reflects the poorly drained nature of the soil.

General landuse versatility ratings for McKerchar soils

Note: The versatility ratings in the table below are indicative of the major limitations for semi-intensive to intensive landuse. These ratings differ from those used in the past in that sustainability factors are incorporated in the classification.

Refer to the Topoclimate district soil map or property soil map to determine which of the soil symbols listed below are applicable, then check the versatility ratings for that symbol in the appropriate table.

McU3 (McKerchar undulating shallow)

Versatility evaluation for soil McU3		
Landuse	Versatility rating	Main limitation
Non-arable horticulture	Limited	Inadequate aeration for sustained periods; restricted rooting depth.
Arable	Limited	Inadequate aeration for sustained periods; risk of short-term waterlogging after heavy rain
Intensive pasture	Limited	Risk of short-term waterlogging after heavy rain
Forestry	Limited	Inadequate aeration for sustained periods; restricted rooting depth.

Management practices that may improve soil versatility

- Careful management after heavy rain and wet periods will reduced the impact of short-term waterlogging. Intensive stocking, cultivation and heavy vehicular traffic use should be minimised during these periods.
- Installation of drainage systems to lower the watertable

Soil profiles available for McKerchar soils

Soil symbol	Profile ID	Topoclimate map sheet	Profile description available	Physical data available	Chemical data available	Profile photo available
McU3	MT10	7	✓	✓	✓	✓

Published by Crops for Southland with financial support from Environment Southland.

Copyright © 2002, Crops for Southland

This Technical Data Sheet may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgement of the source is made. Crops for Southland and Environment Southland would appreciate receiving a copy of any publication that uses this Technical Data Sheet as a source.

No use of this Technical Data Sheet may be made for resale or for any other commercial purpose whatsoever without prior permission in writing from Crops for Southland.

Crops for Southland
PO Box 1306, Invercargill. New Zealand



www.cropssouthland.co.nz